

**“Development of a web-based grape and apple disease risk assessment system”
Progress Report**

**SRSFC Project # 2012 E-02
Extension Proposal**

Name, Mailing and Email Address of Principal Investigator(s):

PRINCIPAL INVESTIGATOR:

Mizuho Nita

Grape Plant Pathologist
AHS Jr. AREC; Virginia Tech
595 Laurel Grove Road
Winchester, VA 22602
Phone: (540) 869-2560
FAX: (540) 869-0862
Email: nita24@vt.edu

Collaborators:

Keith Yoder

Professor
AHS Jr. AREC; Virginia Tech
595 Laurel Grove Road
Winchester, VA 22602
Phone: (540) 869-2560
FAX: (540) 869-0862
Email: ksyoder@vt.edu

Peter Sforza

Director of the Center for Geospatial
Information Technologies
Virginia Tech
113 Major Williams Hall
Blacksburg, VA 24061-0115
Phone: (540) 231-1867
Email: sforza@vt.edu

Seth Peery

Senior GIS Architect
Enterprise GIS Research and
Development Administration
Virginia Tech
Phone: (540) 231-2178
Email: sspeery@vt.edu

Paul Knight

Department of Meteorology
605 Walker Building
University Park, PA 16802
Telephone (814) 863-1842
E-mail: pgk2@psu.edu

Erick De Wolf

Extension Plant Pathologist
Department of Plant Pathology
Kansas State University
4607 Throckmorton Plant Sci. Center
Manhattan, KS 66506
(785) 532-3968
dewolf1@ksu.edu

Objectives:

Establish an online, map-based disease risk assessment system for major fungal and bacterial diseases of grape in Virginia

Justification and Description:

Due to high humidity during the growing season, diseases can have a substantial economic impact on the Virginia fruit industry. Timely application of pesticides is critical for management of many of the fungal and bacterial diseases that affect grape and apple. Often time, growers need to apply preventative pesticides prior to an infection event in order to prevent disease, except a few cases where curative pesticides are available (Ellis 2005). Therefore, growers tend to apply pesticides at regular intervals (i.e., every 7 to 10 days) regardless of weather conditions. However, if weather is dry, many of the fungal and bacterial pathogens are not active, and there is no need for pesticide application for these diseases. Thus, a risk assessment tool that provides specific information about the risk of disease development can be a very useful tool for growers' fungicide decision-making, especially for intensively managed crops

such as grape and apple (Gadoury et al., 1990). What we envision for the future of our program is the establishment of a web-based information center for apple and grape diseases for Virginia growers (Table 1, placed in page 5). We made a progress in 2011 to establish weather data network and developed several candidate grape disease model. In 2012 proposal, we requested a continuation of 2011 to add more candidate disease model, development of a web interface, and also investigate more on weather data retrieval methods.


Development of such a system requires a careful planning and establishment of a good infrastructure so that the system can run efficiently for a long time. Therefore, the entire project was divided into several phases that have specific objectives (Table 1, adjusted from 2011). A portion of the initial two phases of system development was supported by the SRSFC. The key steps during the proposed phase are: 1) the establishment of a data management framework; 2) selection of candidate disease predictive models; 3) development of GIS module/application for visualization of disease risk maps; and 4) implementation of a web-based interactive user interface. The funds were utilized to support student workers who provided support to a) select apple and grape disease models, and b) transfer the model information into GIS modules.

Methodologies: The proposed risk assessment system will have two layers of spatial hierarchy. The top layer will be similar to the currently available Fusarium system where model-estimated weather data will be utilized to create disease risk maps (Fig. 1). The second layer will be based on weather stations (both public and private) located nearby or within a target field. Public weather stations will be visible on the map display (circle icons in Fig. 1), and a user can select a particular weather station to obtain more accurate information. In order to provide more local weather information, the FAA (Federal Aviation Association) and Virginia Cooperative Extension Service's weather stations are added to the system. The website will deliver disease risk analysis based on disease prediction models, and the result of these models will be displayed as a map (Fig. 1). Once a user (a grower) accesses the site, he/she selects a crop, the growth stage of grape, a target disease, and target time window (current or forecast), and a risk map will be generated based on the input. The display will be an easy to understand, color-coded map with green, yellow, and red areas for low, moderate, and high risk of the disease, respectively (Fig. 1). The weather information will be updated daily so that growers can visit the site in the morning prior to make a decision. Also, up to 7-day weather forecast information will be used for future risk assessment, which will give growers "heads-up" for upcoming events.

The website will deliver disease risk analysis based on disease prediction models, and the result of these models will be displayed as a map (Fig. 1). Once a user (a grower) accesses the site, he/she selects a crop, the growth stage of grape, a target disease, and target time window (current or forecast), and a risk map will be generated based on the input. The display will be an easy to understand, color-coded map with green, yellow, and red areas for low, moderate, and high risk of the disease, respectively (Fig. 1). The weather information will be updated daily so that growers can visit the site in the morning prior to make a decision. Also, up to 7-day weather forecast information will be used for future risk assessment, which will give growers "heads-up" for upcoming events.

Progress made in 2011-2012: 1) **Data management and map display:** We obtained weather data from two different sources 1) PSU's climatology lab (Real-Time Mesoscale Analysis or RTMA data, Fig. 3), 2) VCE's mesonet weather station throughout VA (Fig. 2), and 3) Direct retrieval of RTMA data from NOAA (new addition in 2012). The Virginia Tech's Enterprise GIS group who hosts the web application on the Virginia Tech Enterprise GIS system developed a database and network structure where these weather data arrive daily. Then the same

Fig. 1 Mock-up of grape disease risk forecasting system



Impact statements: This project is still under development and it is not yet to make an impact. However, once completed, the impact to both grape and apple producers will be very high. It will help them decide their spray program and ideally, help reducing the use of fungicide, and it can be used as an education tool for extension agents and specialist to discuss about grape and apple disease management.

Table 2. Examples of disease prediction models considered for development of risk assessment models for major grape fungal diseases in Virginia

Crop	Disease	Pathogen	Selected references
GRAPE	Black rot	<i>Guignardia bidwellii</i>	Ellis, M. A., et al. (1986), Funt, R. C., et al. (1990)
	Downy mildew	<i>Plasmopara viticola</i>	Tarr S. A. J. (1972), Strizyk, S. (1983), Madden, L. V. et al. (2000), Lalancette, N., et al. (1988)
	Powdery mildew	<i>Erysiphe necator</i>	Correiar, B. R. (1999), Sall, M. A., (1979)
	Botrytis gray mold	<i>Botrytis cinerea</i>	Broome, J. C., et al. (1995), Bulit, J., et al. (1970), Nair, N. G., and Allen, R. N. (1993), Shtienberg, D., and Elad, Y. (1997)
	Phomopsis cane and leaf spot	<i>Phomopsis viticola</i>	Erincik, O., et al. (2003), Nita, M., et al. (2007)
APPLE	Apple Scab	<i>Venturia inaequalis</i>	Jones ed. Skybit (a commercial service)
	Cedar-apple rust	<i>Gymnosporangium juniperi-virginianae</i>	Biggs (2009, website), Mid-Atlantic Orchard Monitoring Guide
	Quince rusts	<i>Gymnosporangium clavipes</i>	Biggs (2009, website), Mid-Atlantic Orchard Monitoring Guide
	Sooty Blotch and Flyspeck	<i>Peltaster fructicola</i> , <i>Geastrum polystigmatum</i> , <i>Leptodontium elatus</i> , <i>Zygophiala jamaicensis</i>	Brown and Sutton (1995)
	Fire blight	<i>Erwinia amylovora</i>	KTFREC (Maryblyt Website), Dewdney, et al. (2007)

Figure 3 Example of RTMA data stored in VT's CGIS server




Figure 4 Example of black rot model displayed as a map based on RTMA data




Table 1. A timeline of the proposed disease information systems and its impact


Phase	Timeline	Objective		Impact
I Development of the base system A) Disease risk assessment system <i>(In progress)</i>	2009-2012	1	Creating the infrastructure: a) Set up a data server; b) Create a GIS interface; c) Data management (<i>initiated in 2010, expected to complete in early 2011</i>)	- Establish a weather data sharing structure between PSU and VT system - Overhaul of datastream of the VES mesonet
		2	Initial development of grape disease risk assessment tool <i>2009-2011: Selection of candidate models based on previous studies (on-going since 2009 season, several models are translated into ArcGIS)</i> <i>2010-2012: Initial system run: testing and calibration of GIS modules, a hindcast of disease risks with existing datasets (initiated in 2010 season)</i>	
		3	Public deployment of the alpha system of disease maps (<i>a web-interface has been developed in 2011</i>)	
II Initial runs A) Disease risk assessment system B) Grape and Apple disease information center <u>(Sub-sections 1 and 2 are Proposed in this study)</u>	2011-2013	1	Validation of models by comparing model outputs with actual observations in the fields - Obtain accurate observation data from canopies + evaluate leaf wetness measurement methods (models) - Study period: July 2011 to June 2013	- Confirm quality of RTMA data - Development of a better estimate for leafwetness period using available data
		2	Validation of weather station input by comparing with national weather service data and RTMA model results (partly on-going since 2009 season)	
		3	Establishment of the web-interface for grape and apple disease information center (<i>on-going 2011-12</i>)	
		4	2011-2014: Initiation of experiments where participating plots will be using results of risk models to schedule fungicide application	- Student education - Public awareness development - Extension education
		5	Public deployment of the beta system that includes grape and apple disease information such as factsheets and pesticide spray recommendations (existing information will be fully utilized)	
III Public deployment A) Disease risk assessment system B) Grape and Apple disease information center	2012-2015	1	System deployment and feedback: a) System maintenance and evaluation, b) Conduct survey to obtain user comments and suggestions; c) Continuation of validation of models and weather stations	
		2	Use of disease risk assessment system as an extension education tool	
IV Value addition A) Disease risk assessment system B) Grape and Apple disease information center	2011-2016	1	Toward more comprehensive information system for grape management; 1) Implementation of other disease risk maps (Pierce's disease forecast, grape leafroll virus distribution map, etc); 2) Establish links to existing disease databases (e.g., ipmPIPE products) and other information sources 3) Establish apple disease module which is another important fruit crop in VA	- More extended collaboration among faculties and institutions - Extension education beyond mid-Atlantic grape and apple production
		2	Expansion of the system beyond VA and plant pathology: 1) Continuation of the validation of the system; 2) Expansion of the system to other states; 3) Consult viticulturists and entomologists for application of the system; 4) Expansion of the system beyond grape plant pathology.	

Appendix: A poster made for CGIT to describe progress we made so far to general public.

Selected Project: Grape Disease Forecasting

CGIT is creating a predictive model to assess blight risks in grapes. The model considers blossom, canker, shoot, and trauma blight conditions to show infection risks (low to high) to determine risks two days in advance using forecasted weather information.

The model will generate hourly risk prediction rasters to determine when the symptoms of blights would occur. This can help growers to make decisions on the need for blight-preventive measures.



Read RTMA climate data. Accept user inputs for phenology, wetness, and spray conditions.

Calculate Degree Day and Epiphytic Insect Potential.

Do symptoms indicate blight conditions: trauma, shoot, canker and blossom?

NO:
Repeat previous steps.

YES:
1. Calculate probability of blight occurrence and the area of grapes being affected.
2. Assign risk scores
3. Calculate weighted index.