

2014 SRSFC Progress report

Title of Project

Identifying Marketable Attributes in Arkansas Fresh-Market Blackberries

Progress Report

Grant Code

SRSFC Project # 2014-14

Research Proposal

Principal investigators

Principal Investigator:

Dr. Renee Threlfall, Research Scientist, Institute of Food Science and Engineering, **2650 N. Young Ave., University of Arkansas, Fayetteville, AR 72704**, rthrelf@uark.edu

Co-Principal Investigators:

Dr. John R. Clark, University Professor, Department of Horticulture, 316 Plant Science Building, University of Arkansas, Fayetteville, AR 72701, jrclark@uark.edu

Dr. Luke Howard, Professor, Functional Foods Research Program, Department of Food Science, 2650 N. Young Ave., University of Arkansas, Fayetteville, AR 72704, lukeh@uark.edu

Objectives

1. Determine compositional attributes of fresh-market blackberry genotypes from the Arkansas blackberry breeding program
2. Identify nutraceutical attributes of fresh-market blackberry genotypes from the Arkansas blackberry breeding program

Justification

Blackberry (*Rubus subgenus Rubus*) is one of the best examples of a wild-harvested specialty crop that moved to commercial use through breeding efforts. This nutraceutical-rich, fresh-market fruit has the potential for an increased role in commercial markets due to consumers' increasing demand for food products with high functional/health properties. Public and private blackberry breeding programs play a critical role in the future of the blackberry industry. In the South the largest blackberry breeding effort is conducted at the University of Arkansas, Fayetteville (directed by John R. Clark) The Arkansas program contributes the majority of the varieties for the South and is the primary program to help in the southern blackberry industry development. Breeding for enhanced nutraceutical composition of blackberries has been hampered by lack of information on genetic markers that influence genetic and environmental controls and the lack of published information on nutraceutical composition of blackberries. The South has an underutilized capability for the production of fresh-market blackberries and there is a need to identify composition-and nutraceutical-based marketable attributes of fresh-

market blackberry genotypes from the University of Arkansas black berry breeding program. As with all crops, breeding and release of new cultivars to address evolving changes and production challenges is vital to keep healthy markets. Data generated from the proposed work will also provide information used in marketing these genotypes .

Methodology:

The composition and nutraceutical attributes of fresh-market blackberries were addressed. Fruit was harvested at the University of Arkansas Fruit Research Station, Clarksville in June 2014. Four cultivars ('Natchez', 'Osage', 'Ouachita', and 'PrimeArk® 45') and 25 advanced selections were harvested. After harvest, the fruit was taken to the Department of Food Science, University of Arkansas, Fayetteville for evaluation of composition and nutraceutical attributes. The experiment was designed as a completely randomized design. The composition and nutraceutical attributes were or will be evaluated with three replicated samples for each genotype. Analyses will be conducted using JMP® (version 11.0; SAS Institute Inc., Cary, NC). Tukey's HSD (Honestly Significant Difference) will be used for mean separation. Pearson's correlation will be used to test the relationship between/within attributes.

Step 1. Methods for composition analysis (completed June-October 2014)

Three samples of approximately 100 g of berries were collected for each cultivar or genotype, placed in plastic storage bags, and stored at -20°C until analysis. From the frozen berries, three berries per genotype and replication were used to determine berry attributes (individual berry weight, berry length, and berry width) and pyrene attributes (number/berry and dry weight/berry). Three replicate three-berry samples of each cultivar and genotype were used to determine soluble solids, pH, and titratable acidity for each genotype.

Step 2. Methods for nutraceutical analysis (started November 2014 and will be completed by January 2015)

Nutraceutical analysis will be done on each genotype in triplicate beginning in November 2014. To obtain sample extracts, samples (25 g) will be homogenized with 20 mL of acetone/water/acetic (70:29.5:0.5 v/v/v) with a Euro Turrax T18 TissueMizer. The samples will be filtered through Miracloth, the filter cakes will be isolated, and the extraction will be repeated. The filtrates will be adjusted to a final volume of 250 mL with extraction solvent. Samples will be analyzed by HPLC for ellagitannins and flavonols, total monomeric anthocyanins, total phenolics. Oxygen Radical Absorbance Capacity (ORAC) values will be determined on a dual pump BMG Fluostar Optima plate reader.

Results:

Compositional analysis has been completed, but the nutraceutical analysis is in progress. Twenty nine blackberry genotypes were evaluated.

Table 1. Composition attributes for blackberry genotypes Clarksville, AR 2014.

Genotype	Soluble solids (%)	pH	Titrateable acidity (g/L) ^z
A-2252	9.80 cdef ^y	3.58 ab	0.64 d
A-2312	8.83 ef	3.42 abcd	1.02 abcd
A-2316	9.90 cdef	3.10 bcd	1.24 abcd
A-2416	9.47 def	3.16 abcd	1.01 abcd
A-2418	8.07 f	3.08 cd	1.35 ab
A-2419	10.03 cdef	3.06 cd	1.47 a
A-2428	9.87 cdef	3.18 abcd	0.98 abcd
A-2434	9.73 def	3.05 d	1.16 abcd
A-2435	10.33 bcdef	3.33 abcd	0.82 abcd
A-2444	12.30 abc	3.44 abcd	0.67 d
A-2450	8.93 ef	3.00 d	1.16 abcd
A-2452	10.47 bcdef	3.18 abcd	1.33 abc
A-2453	10.63 bcde	3.37 abcd	0.75 bcd
A-2454	10.20 bcdef	3.29 abcd	0.82 abcd
A-2473	10.90 abcde	3.42 abcd	0.90 abcd
A-2480	12.67 ab	3.53 abc	1.00 abcd
A-2487	11.83 abcd	3.61 a	0.79 bcd
A-2491	10.97abcde	3.20 abcd	0.97 abcd
APF-190	8.97 ef	3.20 abcd	0.92 abcd
APF-238	13.33 a	3.25 abcd	0.78 bcd
APF-266	9.13 ef	3.18 abcd	0.84 abcd
APF-268	9.93 cdef	3.16 abcd	0.88 abcd
APF-290	10.17 bcdef	3.29 abcd	1.21 abcd
APF-293	8.67 ef	3.26 abcd	0.97 abcd
APF-298	11.20 abcde	3.26 abcd	1.08 abcd
Natchez	10.17 bcdef	3.17 abcd	1.03 abcd
Osage	8.90 ef	3.58 ab	0.69 cd
Ouachita	10.60 bcdef	3.43 abcd	0.66 d
Prime-Ark45	9.47 def	3.38 abcd	0.81 bcd

^z Titrateable acidity expressed as citric acid

^y Genotypes were evaluated in triplicate (n=3). Means with different letter(s) for each attribute are significantly different (p < 0.05) using Tukey's HSD

Table 3. Berry attributes for blackberry genotypes Clarksville, AR 2014.

Genotype	Berry weight (g)	Berry length (mm)	Berry width (mm)	Berry volume (mm ³) ^z	Drupelet number/ berry
A-2252	6.00 ghijk ^y	28.82 hij	22.15 cdef	3707.58 cde	61.33 ghi
A-2312	8.61 cdefghi	32.88 cdefgh	23.83 abcde	4901.12 bcde	89.00 bcdefg
A-2316	7.13 efghijk	29.89 defghi	21.11 def	3495.84 cde	89.67 bcdefg
A-2416	11.06 bcd	36.80 abcdefg	24.81 abcd	5955.08 abc	88.55 bcdefg
A-2418	9.66 bcde	32.90 cdefgh	25.31 abcd	5562.13 abcd	84.22 cdefgh
A-2419	6.68 efghijk	31.23 cdefghi	21.30 def	3755.66 cde	97.89 abcde
A-2428	8.51 cdefghi	31.34 cdefghi	22.72 abcdef	4237.00 bcde	68.44 fghi
A-2434	9.03 cdefg	34.46 bcdefgh	24.30 abcde	5402.33 abcd	86.22 cdefgh
A-2435	9.19 cdef	36.87 abcdef	24.62 abcde	5859.70 abc	101.89 abcd
A-2444	9.32 cde	30.86 cdefghi	24.11 abcde	4689.53 bcde	52.89 i
A-2450	9.05 cdef	37.32 abcd	22.16 cdef	4844.27 bcde	92.22 bcdef
A-2452	12.66 ab	41.71 ab	26.79 a	7898.96 a	110.44 abc
A-2453	6.01 ghijk	27.47 hij	24.85 abcd	4441.41 bcde	50.22 i
A-2454	8.07 defghij	29.65 defghij	24.69 abcde	4740.69 bcde	64.33 fghi
A-2473	5.78 hijk	29.01 ghij	23.19 abcdef	4112.70 bcde	73.11 efghi
A-2480	5.23 jk	21.86 j	19.42 f	2159.95 e	67.44 fghi
A-2487	4.90 k	23.58 ij	21.42 def	2833.99 de	59.11 hi
A-2491	9.70 bcde	36.76 abcdefg	23.07 abcdef	5185.11 abcd	81.78 defgh
APF-190	8.45 cdefghi	30.80 cdefghi	22.20 bcdef	4010.05 bcde	76.11 defghi
APF-238	5.64 ijk	27.13 hij	22.73 abcdef	3683.11 cde	53.34 i
APF-266	11.12 bc	36.93 abcde	26.08 abc	6600.57 ab	110.33 abc
APF-268	9.07 cdef	36.97 abcde	24.01 abcde	5608.27 abcd	78.22 defghi
APF-290	6.85 efghijk	29.18 efghij	22.31 bcdef	3821.37 bcde	82.56 cdefgh
APF-293	8.97 cdefg	38.36 abc	23.72 abcdef	5675.13 abc	115.67 ab
APF-298	6.27 fghijk	31.60 cdefgh	20.40 ef	3483.46 cde	82.89 cdefgh
Natchez	14.26 a	43.68 a	25.96 abc	7726.93 a	125.83 a
Osage	7.29 efghijk	27.81 hij	26.92 a	5374.68 abcd	70.22 efghi
Ouachita	8.80 cdefgh	29.08 fghij	26.56 ab	5458.53 abcd	69.78 efghi
Prime-Ark45	7.64 efghijk	33.43 cdefgh	22.71 abcdef	4534.55 bcde	85.78 cdefgh

^z Volume calculated as a cone

^y Genotypes were evaluated in triplicate (n=3). Means with different letter(s) for each attribute are significantly different (p < 0.05) using Tukey's HSD

Table 3. Pyrene attributes for blackberry genotypes Clarksville, AR 2014.

Genotype	Pyrene weight (g)/berry	Pyrenes/berry	Pyrene weight/berry weight (%)
A-2252	0.19 fgh ^z	58.44 ijk	3.18 bcd
A-2312	0.27 cdefgh	90.33 cdefg	3.26 abcd
A-2316	0.30 bcdef	91.89 bcdefg	4.21 ab
A-2416	0.28 cdefgh	83.22 defghij	2.52 d
A-2418	0.38 abc	94.22 bcdefg	3.90 abcd
A-2419	0.27 cdefgh	106.89 abcde	4.04 abc
A-2428	0.29 cdefgh	69.11 fghijk	3.35 abcd
A-2434	0.36 abcd	88.78 cdefgh	4.03 abc
A-2435	0.25 defgh	96.11 abcdef	2.76 bcd
A-2444	0.24 efgh	54.00 k	2.61 cd
A-2450	0.32 abcde	92.00 bcdefg	3.58 abcd
A-2452	0.40 ab	111.78 abcd	3.19 bcd
A-2453	0.18 h	51.00 k	2.93 bcd
A-2454	0.25 defgh	68.56 fghijk	3.15 bcd
A-2473	0.23 efgh	76.00 fghijk	3.96 abcd
A-2480	0.20 fgh	70.89 fghijk	3.77 abcd
A-2487	0.19 fgh	59.89 hijk	3.92 abcd
A-2491	0.28 cdefgh	84.78 defghij	2.92 bcd
APF-190	0.27 cdefgh	78.44 efghijk	3.20 bcd
APF-238	0.18 gh	55.89 jk	3.20 bcd
APF-266	0.29 bcdefg	123.33 a	2.63 cd
APF-268	0.29 bcdefg	78.22 efghijk	3.21 abcd
APF-290	0.26 defgh	88.22 cdefgh	3.87 abcd
APF-293	0.42 a	120.66 ab	4.70 a
APF-298	0.21 efgh	86.44 cdefghi	3.38 abcd
Natchez	0.43 a	115.00 abc	3.12 bcd
Osage	0.26 defgh	72.33 fghijk	3.62 abcd
Ouachita	0.23 efgh	66.78 ghijk	2.66 cd
Prime-Ark45	0.32 abcde	90.22 cdefg	4.21 ab

^z Genotypes were evaluated in triplicate (n=3). Means with different letter(s) for each attribute are significantly different (p < 0.05) using Tukey's HSD

Discussion and conclusions:

The group of genotypes in the study provided a substantial range in variables measured. One of the first focus areas was that of soluble solids and pH/titratable acidity, as these play a major role in flavor of blackberries, with the sweeter, lower-acid berries usually preferred by consumers. Selection APF-238 had the highest soluble solids (13.33%) and other values ranged down to A-2418 which had the lowest soluble solids (8.07%). The pH ranged from a high of 3.61 for A-2487 to the low of 3.00 for A-2450. Likewise, the titratable acidity values ranged from 0.64% (A-2252) to 1.47 (A-2419). It was very interesting to note that those berries usually observed to have the more desirable flavor usually had a titratable acidity value under 1.0%. This could be a key indicator in evaluation and breeding for increased flavor and acceptance.

In addition to the soluble solids, pH/titratable acidity, the other physicochemical attributes of blackberries (size, number of drupelets, number of pyrenes) play a key role in acceptance. These other attributes had substantial range in values. 'Natchez' blackberries had the highest berry weight (14.26 g), berry length (43.68 mm), number of drupelets/berry (125.83), and pyrene weight/berry (0.43 g). Whereas, A-2487 had the lowest berry weight (4.90 g), A-2480 had the lowest berry length (21.86 mm), and A-2453 had the lowest number of drupelets/berry (50.22) and the lowest pyrene weight/berry (0.18 g). A-2452 had the highest berry volume (7898.96 mm³) and A-2480 had the lowest (2159.95 mm³). Berry width ranged from 26.92 mm for 'Osage' and 19.42 mm for A-2480. The number of pyrenes/berry ranged from 123.33 (APF-266) to 51.00 (A-2453). APF-293 had the highest pyrene weight/berry weight (4.70%) and A-2416 had the lowest (2.52%).

Impact statement:

Initial results identified blackberry attributes that can be used by the University of Arkansas blackberry breeding program during breeding selection process.

Citations for any publications arising from the project

Not publications or presentations have resulted from this work, but are planned for several meetings in 2015.